

Research Article

The Unique Maximal GF -Regular Submodule of a Module

Areej M. Abduldaïm and Sheng Chen

Department of Mathematics, Harbin Institute of Technology, Harbin 150001, China

Correspondence should be addressed to Areej M. Abduldaïm; areej.mussab@yahoo.com

Received 20 July 2013; Accepted 15 August 2013

Academic Editors: J. He, I. G. Ivanov, J. Rada, and L. Székelyhidi

Copyright © 2013 A. M. Abduldaïm and S. Chen. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

An R -module A is called GF -regular if, for each $a \in A$ and $r \in R$, there exist $t \in R$ and a positive integer n such that $r^n tr^n a = r^n a$. We proved that each unitary R -module A contains a unique maximal GF -regular submodule, which we denoted by $MGF(A)$. Furthermore, the radical properties of A are investigated; we proved that if A is an R -module and K is a submodule of A , then $MGF(K) = K \cap MGF(A)$. Moreover, if A is projective, then $MGF(A)$ is a G -pure submodule of A and $MGF(A) = M(R) \cdot A$.

1. Introduction

Throughout this paper, R is a commutative ring with identity and all modules are left unitary, unless otherwise stated. Recall that an element $r \in R$ is said to be regular if there exists $t \in R$ such that $rtr = r$; a ring R is called regular if and only if each element of R is regular. An ideal I of a ring R is regular if each of its elements is regular in R ; indeed, a regular ideal I of R is itself a regular ring [1]. Brown and McCoy proved in [1] that each ring R contains a unique maximal regular ideal $M(R)$ which satisfies the well-known radical properties. The ideal $M(R)$ is called the regular radical of R .

The concept of regularity was extended to modules in several ways and in [2] the notion of F -regular modules (in the sense of Fieldhouse [3]) was generalized to GF -regular modules. Let A be an R -module; an element $a \in A$ is said to be GF -regular if for each $r \in R$ there exist $t \in R$ and a positive integer n such that $r^n tr^n a = r^n a$. An R -module A is called GF -regular if and only if all its elements are GF -regular; in particular, a ring R is GF -regular if and only if R is GF -regular as an R -module. On the other hand a ring R is π -regular if and only if R is a GF -regular R -module; recall that a ring R is π -regular if, for each $r \in R$, there exist $t \in R$ and a positive integer n such that $r^n tr^n = r^n$. A submodule N of an R -module A is called GF -regular if each element of N is GF -regular and every submodule of a GF -regular module is a GF -regular module. Also, in [2] the concept of G -pure submodules was introduced; a submodule P of an R -module

A is called G -pure if, for each $r \in R$, there exists a positive integer n such that $P \cap Rr^n A = Rr^n P$.

In this paper we show that each module contains a unique maximal GF -regular submodule, which we denote by $MGF(A)$, and we show that $MGF(A)$ satisfies some but not all of the usual radical properties.

2. Main Results

Theorem 1. *Let R be any ring. Every R -module contains a unique maximal GF -regular submodule.*

Proof. Let R be any ring, let A be an R -module, and let

$$G = \{N \mid N \text{ is a } GF\text{-regular submodule of } A\}, \quad (1)$$

where $G \neq \emptyset$ because (0) is a GF -regular submodule of A . Let $\{N_i\}$ be an ascending chain in G and $B = \bigcup_{i \in \Lambda} N_i$. Let $b \in B$; there exists $j \in \Lambda$ such that $b \in N_j$, but N_j is a GF -regular submodule; then, for each $r \in R$, there exist $t \in R$ and a positive integer n such that $r^n tr^n b = r^n b$; therefore b is a GF -regular element in B which implies that B is a GF -regular R -module. Now, by Zorn's lemma, G contains a maximal element which we call MGF . To prove the uniqueness of MGF , assume that MGF_1 and MGF_2 be two maximal GF -regular submodules in A ; then for any maximal ideal P of R each of MGF_1_P and MGF_2_P is semisimple over R_P [2, Proposition 21]. Now, let $MGF_1_P \cap MGF_2_P = K_P$; then $K_P \subseteq MGF_1_P$ and $K_P \subseteq MGF_2_P$; thus $MGF_1_P = K_P + A1_P$ and

$MGF2_p = K_p + A2_p$, where $A1_p$ and $A2_p$ are two submodules of A_p [4]. Hence, $MGF1_p + MGF2_p = A1_p + K_p + A2_p$, but each of $A1_p$, $A2_p$, and K_p is a semisimple submodule; thus $MGF1_p + MGF2_p$ is a semisimple submodule which implies that $MGF1_p + MGF2_p$ is GF -regular [2]. So $MGF1 + MGF2$ is a GF -regular submodule [2, Theorem 20]. Now, each of $MGF1$ and $MGF2$ is a maximal GF -regular submodule and hence $MGF1 + MGF2 = MGF2 = MGF1$. \square

Remark 2. We denote the unique maximal GF -regular submodule of an R -module A by $MGF(A)$. It is obvious that $MGF(A)$ contains every GF -regular submodule of A ; this means that $MGF(A)$ is a GF -regular submodule which is not contained properly in any other GF -regular submodule. In fact, $MGF(A)$ is the sum of all GF -regular submodules of A and $MGF(A) = A$ if and only if A is a GF -regular module.

Example 3. (a) Since the Z -module Z_n is GF -regular for each positive integer n [2], then $MGF(Z_n) = Z_n$.

(b) Each element in the Z -module Q is not GF -regular [2]; hence $MGF(Q) = (0)$.

(c) Let p be a prime number and let $A = Z_{p^\infty} = \bigcup_{i=1}^\infty Z_{p^i}$ be a Z -module. Let $a \in \bigcup_{i=1}^\infty Z_{p^i}$; then there exists a positive integer m such that $a \in Z_{p^m}$, but Z_n is a GF -regular Z -module for each positive integer n ; hence a is a GF -regular element, so $a \in MGF(Z_{p^\infty})$ which implies that $MGF(Z_{p^\infty}) = Z_{p^\infty}$.

(d) Let Q/Z be a Z -module; since Q/Z is a torsion Z -module, then Q/Z is a GF -regular Z -module [2, Proposition 6]. Since $Q/Z = \sum_p Z_{p^\infty}$ for each prime number p , then $MGF(Q/Z) = \sum_p Z_{p^\infty}$ for all primes p .

Proposition 4. Let A and B be R -modules, and let K be a submodule of A ; then

$$(a) \ MGF(K) = K \cap MGF(A),$$

$$(b) \ MGF(A \oplus B) \subseteq MGF(A) \oplus MGF(B).$$

Proof. (a) Let K be a submodule of A , and let $k \in MGF(K)$; then $k \in K$ and k is GF -regular in K which implies that k is GF -regular in A , thus $k \in K \cap MGF(A)$. Conversely, let $k \in K$ and $k \in MGF(A)$; therefore k is GF -regular in K which means that $k \in MGF(K)$ and hence $MGF(K) = K \cap MGF(A)$.

(b) Let $c \in MGF(A \oplus B)$; then $c = (a, b)$, where $a \in A$ and $b \in B$. Since c is GF -regular, then each of a and b is GF -regular which means that $a \in MGF(A)$ and $b \in MGF(B)$; hence $c \in MGF(A) \oplus MGF(B)$. \square

Proposition 5. Let A and A' be R -modules, and let $f : A \rightarrow A'$ be an R -homomorphism; then $f(MGF(A)) \subseteq MGF(f(A))$.

Proof. If $a \in MGF(A)$, then $R/\text{ann}(a)$ is a π -regular ring, but $\text{ann}(a) \subseteq \text{ann}(f(a))$; thus $R/\text{ann}(f(a))$ is an epimorphic image of $R/\text{ann}(a)$; hence it is a π -regular ring. Therefore, $f(a) \in MGF(f(A))$ and $f(MGF(A)) \subseteq MGF(f(A))$. \square

Remark 6. (a) If $f : A \rightarrow A'$ is an R -epimorphism, then $f(MGF(A)) \neq MGF(A')$ in general. In fact, let $\pi : Z \rightarrow Z_4 \cong Z/4Z$ be the natural map, where Z and Z_4 are Z -modules. It

is easy to check that $MGF(Z) = (0)$, $f(MGF(Z)) = (0)$, but $MGF(Z_4) \cong Z_2$.

(b) It is shown in [1] that, for a ring R , $M(R/M(R)) = (0)$ which is not true in case of GF -regular modules; this means that $MGF(A/MGF(A)) \neq (0)$ (as in (a)).

Corollary 7. For each R -module A , $M(R) \cdot A \subseteq MGF(A)$.

Proof. For each $a \in A$, let $f : R \rightarrow A$ be an R -homomorphism defined by $f(r) = ra$. Then $f(M(R)) \subseteq MGF(A)$ by Proposition 5, but $M(R) \cdot A = \sum_a f(M(R))$; hence $M(R) \cdot A \subseteq MGF(A)$. \square

Let $J(R)$ be the Jacobson radical of a ring R . Brown and McCoy proved in [1] that $M(R) \cap J(R) = (0)$. However, this is not true for GF -regular modules; for example, if $A = Z_4$ is a Z -module, then $MGF(A) \cong Z_2$, $J(A) \cong Z_2$ and $MGF(A) \cap J(A) \neq (0)$.

Lemma 8. Let A be an R -module and let P be a G -pure submodule of A . For any $r \in R$, there exists a positive integer n such that $P = Rr^n P$ if and only if $P \subseteq Rr^n A$.

Proof. Since P is G -pure in A , then, for each $r \in R$, there exists a positive integer n such that $P \cap Rr^n A = Rr^n P$. If $P = Rr^n P$, then $P \cap Rr^n A = P$, and hence $P \subseteq Rr^n A$. Conversely, if $P \subseteq Rr^n A$, then $P \cap Rr^n A = P$, but $P \cap Rr^n A = Rr^n P$; therefore $P = Rr^n P$. \square

Lemma 9. Let $r \in J(R)$; if P is a finitely generated G -pure submodule of an R -module A such that $P \subseteq Rr^n A$ for some positive integer n , then $P = 0$.

Proof. By Lemma 8 we get that $P = Rr^n P$ and by Nakayama's lemma [5], $P = 0$. \square

Theorem 10. Let A be an R -module. If $MGF(A)$ is a G -pure submodule of A , then $MGF(A) \cap J(R) \cdot A = (0)$.

Proof. Let $r \in MGF(A) \cap J(R) \cdot A$, and let $P = Rr$. It is clear that $P \subseteq MGF(A)$. Since $MGF(A)$ is a GF -regular module, then P is a G -pure submodule in $MGF(A)$ [2, Theorem 11]. But $MGF(A)$ is G -pure in A ; hence P is G -pure in A . Now, $P \subseteq J(R) \cdot A$, so $P = 0$ by Lemma 9. Therefore $MGF(A) \cap J(R) \cdot A = (0)$. \square

Recall that $M(R)$ is always a pure ideal in R . Hence $M(R)$ is G -pure [2].

Theorem 11. Let A be a projective R -module; then

$$(a) \ MGF(A) = M(R) \cdot A,$$

$$(b) \ MGF(A) \text{ is a } G\text{-pure submodule of } A,$$

$$(c) \ MGF(A) \cap J(A) = (0).$$

Proof. (a) By the dual basis lemma [4], for each $a \in A$ we have that $a = \sum_i f_i(a)a_i$, where $a_i \in A$ for all i and $f_i \in A^* := \text{Hom}_R(A, R)$. If $a \in MGF(A)$, then the submodule Ra is GF -regular and $f_i(Ra)$ is a GF -regular ideal in R by Proposition 5, hence $M(R)$. Thus $MGF(A) \subseteq M(R) \cdot A$. We get the other direction of the inclusion by Corollary 7.

(b) First we claim that; for any two ideals K and L of R , $(K \cap L)A = KA \cap LA$; it is enough to show this locally; thus we may assume that A is free. It is clear that $(K \cap L)A \subseteq KA \cap LA$. On the other hand, let $x \in KA \cap LA$; then

$$x = \sum r_i x_i = \sum s_i x_i \quad r_i \in K, \quad s_i \in L. \quad (2)$$

By freeness, $r_i = s_i$ and $x \in (K \cap L)A$. Now, let K be any ideal in R ; then by (a) we get $MGF(A) \cap KA = M(R) \cap KA = (M(R) \cap K)A$. But $M(R)$ is G -pure ideal, so, for each $r \in R$, there exists a positive integer n such that $M(R) \cap Rr^n = Rr^n M(R)$; hence $MGF(A) \cap Rr^n A = Rr^n M(R) \cdot A = Rr^n MGF(A)$.

(c) Since A is projective, then $J(A) = J(R) \cdot A$ [4] which implies that $MGF(A) \cap J(A) = (0)$ by Theorem 10. \square

Corollary 12. Let R be any ring, and let A be any R -module such that $MGF(A)$ is a G -pure submodule and $J(A) = J(R) \cdot A$; then $MGF(A) \cap J(A) = (0)$.

Proof. Since $J(A) = J(R) \cdot A$, then by Theorem 10 we get that $MGF(A) \cap J(A) = (0)$. \square

Remark 13. If A is a GF -regular R -module, then $MGF(A) = A$; hence $J(R) \cdot A = J(R) \cdot A \cap A = J(R) \cdot A \cap MGF(A) = (0)$. In fact, this shows that Theorem 10 is a generalization of [2, Proposition 28].

In [2] we noticed that every module over π -regular ring is GF -regular, but the converse need not be true in general. The next result shows how the converse may be true, but first we recall that if A is an R -module, then the trace of A is $\text{tr}(A) = \sum_{f \in A^*} f(A)$, where $A^* = \text{Hom}(A, R)$.

Proposition 14. Let A be a GF -regular R -module. If $\text{tr}(A) = R$, then R is a π -regular ring.

Proof. For each $a \in A$ and $f \in A^* = \text{Hom}(A, R)$, since Ra is a GF -regular submodule of A , then by [2, Proposition 7] we get that $f(Ra)$ is a π -regular ideal. Thus $f(Ra) \subseteq M(R)$, but $f(Ra) \subseteq \text{tr}(A)$; hence $\text{tr}(A) = R \subseteq M(R)$, which implies that $M(R) = R$ and R is π -regular. \square

Conflict of Interests

The authors declare that there is no conflict of interests regarding the publication of this paper.

Acknowledgments

Sheng Chen was supported by the National Natural Science Foundation of China (Grants nos. 11001064 and 11101105), by the Fundamental Research Funds for the Central Universities (Grant no. HIT. NSRIF. 2014085), and by the Scientific Research Foundation for the Returned Overseas Chinese Scholars, State Education Ministry, and Areej M. Abduldaim was supported by China Scholarship Council (CSC no. 2011368T09).

References

- [1] B. Brown and N. H. McCoy, "The maximal regular ideal of a ring," *Proceedings of the American Mathematical Society*, vol. 1, pp. 165–171, 1950.
- [2] A. M. Abduldaim and S. Chen, "GF-regular modules," *Journal of Applied Mathematics*, vol. 2013, Article ID 630285, 7 pages, 2013.
- [3] D. J. Fieldhouse, *Purity and flatness [Ph.D. thesis]*, McGill University, Montreal, Canada, 1967.
- [4] F. Kacsh, *Modules and Rings*, vol. 17, Academic Press, New York, NY, USA, 1982.
- [5] M. D. Larsen and P. J. MacCarthy, *Multiplicative Theory of Ideals*, Academic Press, New York, NY, USA, 1971.

